JOURNAL OF PALEONTOLOGY

INDEX TO VOLUME 28

AUTHOR INDEX

		No.	Page
1.	AKERS, W. H., Ecologic aspects and stratigraphic significance of the foraminifer *Cyclammina cancellata* Brady	2	132
2.	ANISGARD, HARRY W., Echinoid from the Eocene of western Venezuela, An	6	830
3.	ARKELL, W. J., Nomenclature of families and superfamilies	2	218
4.	ARNOLD, ZACH M., Culture methods in the study of living Foraminifera	4	404
5.	AVNIMELECH, M., PARNESS, A., and REISS, Z., Mollusca and Foraminifera from the lower Albian of the Negev (southern Israel)	6	835
	BISHOP, ERNEST W., see SCHROEDER, MELVIN C. (74)		
6.	BISWAS, BUDDHADEB, On the occurrence of *Hantkenina alabamensis* from the Khasi Hills, Assam, India	6	791
7.	BOCK, WILHELM, *Primaraucaria*, a new araucarian genus from the Virginia Triassic	1	32
8.	BOWSHER, ARTHUR L., Stratigraphic significance of a crinoid from the Redwall limestone of Arizona, The	1	113
9.	BRAMSLETTE, M. N., and RIEDEL, W. R., Stratigraphic value of discoasters and some other microfossils related to Recent coccolithophores	4	385
10.	BRÜCKNER, WERNER D., Note on some fossils from the Girón group in the Rio Lebrija Valley, Department of Santander, Colombia	1	112
11.	BURK, C. A., Faunas and age of the Amsden formation in Wyoming	1	1
12.	CHAFFEE, ROBERT G., *Campylocynodon personi*, a new Oligocene carnivore from the Beaver Divide, Wyoming	1	43
13.	CHEETHAM, ALAN H., New early Cretaceous cheilostome bryozoan from Texas, A	2	177
14.	COLEY, TYROL B., Stratigraphic distribution and correlation of some Middle Devonian Ostracoda	4	452
15.	COOPER, G. ARTHUR, Unusual Devonian brachiopods	3	325
16.	COX, L. R., Lower Cretaceous Mollusca from Pointe-à-Pierre, Trinidad	5	622
17.	DECKER, CHARLES E., New graptolite from South Dakota, A	2	208
	DECKER, JACK M., see KUMMEL, BERNHARD (41)		
18.	DUNBAR, CARL O., New technique for making sections of invertebrate fossils, A	1	112
19.	DURHAM, J. WYATT, New family of clypeastroid echinoids, A	5	677
20.	ELIAS, MAXIM K., *Cambroporella* and *Coeloclema*, Lower Cambrian and Ordovician bryozoans	1	52
21.	FRITZ, MADELEINE A., *Rhopalonaria lambtonensis*, n. sp.	1	118
	FURNISH, W. M., see MILLER, A. K. (47)		
22.	GUTSCHICK, R. C., Holothurian sclerites from the Middle Ordovician of northern Illinois	6	827
23.	GUTSCHICK, R. C., New species of *Astraeospongia* from the Middle Ordovician of northern Illinois, A	4	430
24.	HAAG, WILLIAM G., Proceedings of the First Pan-African Congress on Prehistory, L. S. B. Leakey and Sonia Cole, editors (Review of)	6	874
25.	HAAG, WILLIAM G., The tree of human history by Alan Houghton Brodrick (Review of)	6	874
	HARDING, J. P., see SYLVESTER-BRADLEY, P. C. (82)		
26.	HILL, BERNARD L., Reclassification of winged *Cythereis* and winged *Brachycythere*	6	804
27.	HINTZE, LEHI F., *Presbynileus* and *Protopresbynileus*, new generic names proposed for *Pseudonileus* and *Paranileus* Hintze, preoccupied	1	119
28.	HOWE, HENRY V., Genotype of the ostracod *Moorea*	2	214
29.	HUDSON, R. G. S., Jurassic stromatoporoids from Lebanon	5	657
30.	HUDSON, R. G. S., New Lower Cretaceous stromatoporoid, *Bekhmeia wetzeli*, from northern Iraq, A	1	47
31.	IMLAY, RALPH W., Barremian ammonites from Trinidad, B.W.I.	5	662
32.	JOHNSON, J. HARLAN, Cretaceous Dasycladaceae from Gillespie County, Texas	6	787
	JONES, B. C., see RIVEROLL, D. D. (71)		
33.	KEEN, A. MYRA, *Ventricolaria* and *similivenus insolida*, new names in the pelecypod family Veneridae	2	217
34.	KELLEY, DANA R., and WOOD, ALBERT E., Eocene mammals from the Lysite member, Wind River formation of Wyoming, The	3	337

35. KESLING, ROBERT V., Oncotechmoninae, a new subfamily of entomoconchid ostracods from the Middle Devonian of New York 5 575
36. KING, PHILIP B., The Permian reef complex of the Guadalupe Mountains region, Texas and New Mexico; A study in paleoecology by Norman D. Newell, J. Keith Rigby, Alfred G. Fischer, A. J. Whiteman, John E. Hickox, and John S. Bradley (Review of) 4 493
37. KJELLESVIG-WAERING, ERIK N., Note on a new Silurian (Downtonian) scorpion from Shropshire, England 4 485
38. KOENIG, JOHN W., Application of cellulose peel technique to fenestrate sections 1 76
39. KORNICKER, LOUIS S., Permian *Ancistrum* (Holothuroidea) from Kansas, A 1 117
40. KUMMEL, BERNHARD, Jurassic nautiloids from western North America 3 320
41. KUMMEL, BERNHARD, and DECKER, JACK M., Lower Turonian ammonites from Texas and Mexico 3 310
42. KURTÉN, BJÖRN, Population dynamics—a new method in paleontology 3 286
43. MACDONALD, J. R., New *Pseudaelurus* from the lower Snake Creek fauna of Nebraska, A 1 67
44. MACNEIL, F. STEARNS, *Stenzelia*, new name for *Trinacriella* MacNeil 2 217
45. MAXWELL, W. G. H., *Strophalosia* in the Permian of Queensland 5 533
McFARLAND, HERDIS B., see PECK, JOSEPH H., JR. (61)
46. MILLER, A. K., American Triassic nautiloids, a study of the classification and evolution of late Paleozoic and early Mesozoic cephalopods, and descriptions of new species by Bernhard Kummel (Review of) 4 491
47. MILLER, A. K., and FURNISH, W. M., Classification of the Paleozoic ammonoids, The 5 685
48. MOORE, RAYMOND C., Kingdom of organisms named Protista 5 588
49. MORRIS, WILLIAM J., Eocene fauna from the Cathedral Bluffs tongue of the Washakie Basin, Wyoming, An 2 195
50. MURRAY, GROVER E., Evolution and human destiny by Fred Kohler (Review of) ... 4 491
51. MURRAY, GROVER E., Fossil plants of the Florissant beds, Colorado by Harry D. MacGinitie (Review of) 4 492
52. NICOL, DAVID, Growth and decline of populations and the distribution of marine pelecypods 1 22
53. NICOL, DAVID, Nomenclatural review of genera and subgenera of Cucullaeidae 1 96
54. NORTON, PETER, New Blancan vertebrate locality, A 4 487
55. OKULITCH, VLADIMIR J., Archaeocyatha from the Lower Cambrian of Inyo County, California 3 293
ORMSBEE, JOHN B. S., see WOOD, ALBERT E. (93)
56. PALMER, ALLISON R., Faunas of the Riley formation in central Texas, The 6 709
57. PARKINSON, DONALD, Quantitative studies of brachiopods from the Lower Carboniferous reef limestones of England. I. *Schizophoria resupinata* (Martin) 3 367
58. PARKINSON, DONALD, Quantitative studies of brachiopods from the Lower Carboniferous reef limestones of England. II. *Pugnax pugnus* (Martin) and *P. pseudopugnus* n. sp. 5 563
59. PARKINSON, DONALD, Quantitative studies of brachiopods from the Lower Carboniferous reef limestones of England. III. *Pugnax uminatus* (J. Sowerby) and *P. mesogonus* (Phillips) 5 668
PARNESS, A., see AVNIMELECH, M. (5)
60. PEABODY, FRANK E., Trackways of an ambystomid salamander from the Paleocene of Montana 1 79
61. PECK, JOSEPH H., JR., and McFARLAND, HERDIS B., Whitfield collection types at the University of California 3 297
62. PEDERSON, SELMER L., Permian (Wolfcampian) fauna of the Casper formation of southeastern Wyoming, A 1 17
63. PETERSON, JAMES A., Jurassic Ostracoda from the "lower Sundance" and Rierdon formations, western interior United States 2 153
64. PURI, HARBANS SINGH, Check list of Indian Tertiary larger Foraminifera 2 185
65. RASETTI, FRANCO, Early Ordovician trilobite faunules from Quebec and Newfoundland 5 581
66. RASETTI, FRANCO, Internal shell structures in the Middle Cambrian gastropod *Scenella* and the problematic genus *Stenothecoides* 1 59
67. RASETTI, FRANCO, Phylogeny of the Cambrian trilobite family Catillicephalidae and the ontogeny of *Welleraspis* 5 599
68. REED, CHARLES A., Some fossorial mammals from the Tertiary of western North America 1 102
REISS, Z., see AVNIMELECH, M. (5)
69. RHODES, F. H. T., Value of a diagnosis in systematic paleontology, The 4 487
70. RICHARDS, HORACE G., Analysis of the methods of storing invertebrate fossils, An .. 4 486
RIEDEL, W. R., see BRAMLETTE, M. N. (9)
71. RIVERSOLL, D. D., and JONES, B. C., Varves and Foraminifera of a portion of the upper Puente formation (upper Miocene), Puente, California 2 121

72. RUTSCH, R. F., and SALVADOR, A., Mollusks from the Cogollo and La Luna formations (Cretaceous) of the Chejendé Area, Western Venezuela 4 417
73. SABROSKY, CURTIS W., Nomenclature of families and superfamilies 4 489
SALVADOR, A., see RUTSCH, R. F. (72)
74. SCHROEDER, MELVIN C., and BISHOP, ERNEST W., Notes on Foraminifera from the late Cenozoic in southern Florida 2 210
75. SKINNER, JOHN W., and WILDE, GARNER L., Fusulinid subfamily Boultoniinae, The. 4 434
76. SKINNER, JOHN W., and WILDE, GARNER L., Fusulinid wall structure 4 445
77. SKINNER, JOHN W., and WILDE, GARNER L., New early Pennsylvanian fusulinids from Texas 6 796
78. SMITH, MARVIN L., Method of selecting sample sizes, A 1 116
79. STERNBERG, C. M., Classification of American duck-bulled dinosaurs 3 382
80. STRIMPLE, HARRELL L., New species of *Plummericrinus* 2 204
81. SYLVESTER-BRADLEY, P. C., Form-genera in paleontology 3 333
82. SYLVESTER-BRADLEY, P. C., and HARDING, J. P., Postscript notes on the ostracode *Trachyleberis* 5 560
83. TEICHERT, CURT, New Permian crinoid from Western Australia, A 1 70
84. THALMANN, HANS E., Bibliography and index to new genera, species and varieties of Foraminifera for the year 1953 ..... 6 840
85. TOOMEY, DONALD F., Bibliography of the family Fusulinidae, A 4 465
86. UNKLESBAY, A. G., Distribution of American Pennsylvanian cephalopods 1 84
87. UNKLESBAY, A. G., Nautiloids from the Tanyard formation of central Texas 5 637
88. WALTER, J. C., JR., Correction 1 119
89. WEISS, MALCOLM P., Notes on some Middle Ordovician fossils from Minnesota 4 427
90. WHITTINGTON, HARRY B., New Ordovician graptolite from Oklahoma, A 5 613
WILDE, GARNER L., see SKINNER, JOHN W. (75, 76, 77)
91. WILIMOVSKY, NORMAN J., *Gunnellodus*, a new name for *Idiacanthus* Gunnell 5 693
92. WILSON, JAMES LEE, Late Cambrian and Early Ordovician trilobites from the Marathon Uplift, Texas 3 249
93. WOOD, ALBERT E., and ORMSBEE, JOHN B. S., Notes on mammals from the Upper Cretaceous Lance formation of Wyoming 1 26
WOOD, ALBERT E., see KELLEY, DANA R. (34)

SUBJECT INDEX

Alaska: nautiloids, Jurassic (40).
Albian (Cretaceous): mollusks and foraminifera, Israel (5).
algae: dasyclads, Cretaceous, Texas (32).
ammonites: Barremian (Cretaceous), Trinidad (31); classification, Paleozoic (47); Turonian (Cretaceous), Texas, Mexico (41).
Amsden formation: fauna and age, Wyoming (11).
annelids: possible, Ordovician, Minnesota (89).
Ancistrum (holothurian): Permian, Kansas (39).
Archaeocyatha (pleosponges): Cambrian, California (55).
Arizona: crinoid, Redwall limestone (Mississippian) (8).
Astraeospongia (sponge): Ordovician, Illinois (23).
Australia: crinoid, Permian (83): (Queensland) brachiopods, Permian (45).
Barremian (Cretaceous): ammonites, Trinidad (31).
Bekhmeia wetzeli (stromatoporoid): Cretaceous, Iraq (30).
Boultoniinae (fusulinids): Permian (75).
brachiopods: fauna and age, Amsden formation, Wyoming (11); Permian, Wyoming (62); *Pugnax acuminatus* and *P. mesogonus*, Carboniferous, England (59); *Pugnax pugnus* and *P. pseudopugnus*, Carboniferous, England (58); Riley formation (Cambrian), Texas (56); *Schizophoria resupinata*, Carboniferous, England (58); *Strophalosia*, Permian, Australia (Queensland) (48); unusual, Devonian (15).
Brachycythere (ostracode): Tertiary, Gulf Coast (26).
British Columbia: gastropod, Cambrian (66).
bryozoans: Cambrian and Ordovician (20); Cretaceous, Texas (13); Devonian, Ontario (21).
California: pleosponges, Cambrian (55); varves, foraminifers, Puente formation (Miocene) (71).
Cambrian: bryozoans (20); Catillicephalidae and *Welleraspis* (trilobites) (67); gastropod, British Columbia (66); mollusk? (66); pleosponges, California (55); Riley formation, trilobites, brachiopods, Texas (56); trilobites, Texas (92).
Campylocynodon personi (carnivore): Oligocene, Wyoming (12).
Carboniferous: brachiopods, England (57) (58) (59).
carnivores: *Campylocynodon personi*, Oligocene, Wyoming (12).
Catillicephalidae (trilobites): Cambrian (67).
cellulose peel: technique of application to fenestrate sections (38).

Cenozoic: foraminifers, Florida (74).
cephalopods: Pennsylvanian, distribution in America (86).
coccolithophores: discoasters and other related microfossils, Recent (9).
collections: Whitfield, University of California (61).
Colombia: Girón group, age, fossils (10).
conifer: *Primaraucaria*, Triassic, Virginia (7).
correction: (88).
Cretaceous: ammonites, Trinidad (31); ammonites, Turonian, Texas, Mexico (41); bryozoan, Texas (13); dasyclads (algae), Texas (32); mammals, Lance formation, Wyoming (93); mollusks, foraminifers, Albian, Israel (5); nollusks, Trinidad (16); mollusks, Venezuela (72); stromatoporoid, Iraq (30).
crinoids: Permian, Australia (83); *Plummericrinus*, Pennsylvanian, Oklahoma (80); Redwall limestone (Mississippian), Arizona (8).
Cucullaeidae (pelecypods): nomenclatural review (53).
Cyclammina cancellata (foraminifer): ecologic and stratigraphic aspects (1).
Cythereis (ostracode): Tertiary, Gulf Coast (26).
dasyclads (algae): Cretaceous, Texas (32).
Devonian: bryozoans, Ontario (21); unusual brachiopods (15); ostracodes, Ohio, New York, Ontario (14); new family ostracodes, New York (35).
dinosaurs: American duck-billed, classification (79).
discoasters: microfossils related to Recent coccolithophores (9).
echinoids: Eocene, Venezuela (2); clypeastroid (19).
ecology: *Cyclammina cancellata* (foraminifer) (1).
England: brachiopods, Carboniferous (57) (58) (59): scorpion, Silurian (37).
Eocene: echinoid, Venezuela (2); mammals, Wind River formation, Wyoming (34); mammals, Wyoming (49).
felids: *Pseudaelurus*, Miocene, Nebraska (43).
fish: *Gunnellodus*, new name for *Idiacanthus* (91).
Florida: foraminifers, Cenozoic (74).
foraminifers: Albian (Cretaceous), Israel (5); bibliography of family Fusulinidae (85); Boultoniinae (fusulinids), Permian (75); Cenozoic, Florida (74); *Cyclammina cancellata*, ecologic and stratigraphic aspects (1); culture of living (4); fusulinid wall structure (76); fusulinids, Pennsylvanian, Texas (77); *Hantkenina alabamensis*, India (6); new genera, species, varieties, bibliography and index for 1953 (84); Puente formation (Miocene), California (71); Tertiary, India (64).
Fusulinidae (foraminifers): bibliography (85).
fusulinids (foraminifers): bibliography of family Fusulinidae (85); Boultoniinae, Permian (75); Pennsylvanian, Texas (77); wall structure (76).
gastropods: *Scenella*, Cambrian, British Columbia (66).
Girón group: fossils, age?, Colombia (10).
graptolites: Mississippian, South Dakota (17); Ordovician, Oklahoma (90).
Gulf Coast: ostracodes, Tertiary (26).
Gunnelodus (fish): new name for *Idiacanthus* (91).
Hantkenina alabamensis (foraminifer): India (6).
holothurians: *Ancistrum*, Permian, Kansas (39); Ordovician, Minnesota (89); sclerites, Ordovician, Illinois (22).
Idiacanthus (fish); *Gunnellodus*, new name (91).
Illinois: holothurian sclerites, Ordovician (22); sponge, Ordovician (23).
India: foraminifers, Tertiary (64); *Hantkenina alabamensis* (foraminifer) (6).
invertebrates: storage methods (70); technique for making sections (18).
Iraq: stromatoporoid, Cretaceous (30).
Israel: mollusks and foraminifers, Albian (Cretaceous) (5).
Jurassic: ostracodes, Montana, South Dakota, Wyoming (63); nautiloids, Montana, Oregon, Alaska (40); stromatoporoids, Lebanon (29).
Kansas: holothurian, Permian (39).
Lance formation (Cretaceous): mammals, Wyoming (93).
Lebanon: stromatoporoids, Jurassic (29).
mammals: Eocene, Wyoming (49); Lance formation (Cretaceous), Wyoming (93); Tertiary, Montana, South Dakota (68); Wind River formation (Eocene), Wyoming (34).
Mexico: ammonites, Turonian (Cretaceous) (41).
Minnesota: holothurian, receptaculitid, annelid?, Ordovician (89).
Miocene: felid, Nebraska (43); Puente formation, varves, foraminifers, California (71).
Mississippian: Amsden formation (?), Wyoming (11); crinoids, Redwall limestone, Arizona (8); graptolite, South Dakota (17).
mollusks: Albian (Cretaceous), Israel (5); Cretaceous, Trinidad (16); Cretaceous, Venezuela (72); *Stenothecoides* (?), Cambrian (66); *Stenzelia*, new name for *Trinacriella* (44).
Montana: mammals, Tertiary (68); nautiloids, Jurassic (40); ostracodes, Jurassic (63); salamander tracks, Paleocene (60).
Moorea (ostracode): genotype of (28).

nautiloids: Jurassic, Montana, Oregon, Alaska (40); Ordovician, Texas (87).
Nebraska: felid, Miocene (43).
Newfoundland: trilobites, Ordovician (65).
New York: ostracodes, Devonian (14); new family ostracodes, Devonian (35).
nomenclature: families and superfamilies (3, 73); review of Cucullaeidae (pelecypods) (53).
Ohio: ostracodes, Devonian (14).
Oklahoma: crinoid, Pennsylvanian (80); graptolites, Ordovician (90).
Oligocene: *Campylocynodon personi* (carnivore), Wyoming (12).
Oncotechmoninae (ostracodes): new family, Devonian, New York (35).
Ontario: bryozoan, Devonian (21); ostracodes, Devonian (14).
Ordovician: bryozoans (20); graptolites, Oklahoma (90); holothurian, receptaculitid, annelid?, Minnesota (89); holothurian sclerites, Illinois (22); nautiloids, Tanyard formation, Texas (87); sponge, Illinois (23); trilobites, Quebec, Newfoundland (65); trilobites, Texas (92).
Oregon: nautiloids, Jurassic (40).
ostracodes: *Cythereis* and *Brachycythere*, Tertiary, Gulf Coast (26); Devonian, Ohio, New York, Ontario (14); Jurassic, Montana, South Dakota, Wyoming (63); *Moorea*, genotype of (28); new family, Oncotechmoninae, Devonian, New York (35); notes on *Trachyleberis* (82).
Paleocene: salamander tracks, Montana (60).
paleontology: form-genera (81); methods of storing invertebrate fossils (70); value of diagnosis (69).
Paleozoic: ammonoids, classification (47).
Paranileus (trilobite): new name for (27).
pelecypods: Cucullaeidae, nomenclatural review (53); marine, growth and decline of population (?); *Ventricolaria* and *Similivenus insolida*, new names (33).
Pennsylvanian: Amsden formation (?), Wyoming (11); cephalopods, distribution in America (86); crinoid, Oklahoma (80); fusulinids, Texas (77).
Permian: Boultoniinae (fusulinids) (75); brachiopods, Australia (Queensland) (45); brachiopods, Wyoming (62); crinoids, Australia (83); holothurian, Kansas (39).
Permo-Carboniferous: Girón group (?), fossils, Colombia (10).
pleosponges: Archaeocyatha, Cambrian, California (55).
Plummericrinus (crinoids): Pennsylvanian, Oklahoma (80).
population dynamics: new method in paleontology (42).
Primaraucaria (conifer): Triassic, Virginia (7).
Protista: (48).
Pseudaelurus (felid): Miocene, Nebraska (43).
Pseudonileus (trilobite): new name for (27).
Puente formation (Miocene): varves, foraminifers, California (71).
Pugnax (brachiopod): *acuminatus*, *mesogonus*, Carboniferous, England (59); *pseudopugnus*, *pugnus*, Carboniferous England (58).
Quebec: trilobites, Ordovician (65).
Recent: discoasters and other microfossils related to coccolithophores (9).
receptaculitids: Ordovician, Minnesota (89).
Redwall limestone (Mississippian): crinoid, Arizona (8).
Rhopalonaria lambtonensis (bryozoan): Devonian, Ontario (21).
Riley formation (Cambrian): trilobites, brachiopods, miscellaneous fauna, Texas (56).
salamanders: tracks, Paleocene, Montana (60).
Scenella (gastropod): Cambrian, British Columbia (66).
Schizophoria resupinata (brachiopod): Carboniferous, England (58).
sclerites (holothurian): Ordovician, Illinois (22).
scorpion: Silurian, England (37).
Silurian: scorpion, England (37).
Similivenus insolida (pelecypod): new name (33).
South Dakota: graptolite, Mississippian (17); mammals, Tertiary (68); ostracodes, Jurassic (63).
sponges: *Astraeospongia*, Ordovician, Illinois (23).
Stenothecoides (mollusk?): Cambrian (66).
Stenzelia (mollusk): new name for *Trinacriella* (44).
stromatoporoids: Cretaceous, Iraq (30); Jurassic, Lebanon (29).
Strophalosia (brachiopod): Permian, Australia (Queensland) (45).
Tanyard formation (Ordovician): nautiloids, Texas (87).
techniques: cellulose peel, fenestrate sections (38); invertebrate sections (18); population dynamics (42); selecting sample sizes (78); storage of invertebrates (70).
Tertiary: foraminifers, India (64); mammals, Montana, South Dakota (68); ostracodes, Gulf Coast (26).
Texas: ammonites, Turonian (Cretaceous) (41); bryozoan, Cretaceous (13); dasyclads (algae), Cretaceous (32); fusulinids, Pennsylvanian (77); nautiloids, Ordovician (87); trilobites, brachiopods, miscellaneous fossils, Riley formation (Cambrian), Texas (56); trilobites, Cambrian, Ordovician (92).
Trachyleberis (ostracode): notes on (82).
Triassic: Girón group (?), fossils, Colombia (10); conifer, Virginia (7).

trilobites: Cambrian, Ordovician, Texas (92); Catillicephalidae, *Welleraspis*, Cambrian (67); Ordovician, Quebec, Newfoundland (65); *Pseudonileus* and *Paranileus*, new names for (27); Riley formation (Cambrian), Texas (56).
Trinacriella (mollusk); *Stenzelia*, new name (44).
Trinidad: ammonites, Barremian (Cretaceous) (31); mollusks, Cretaceous (16).
Turonian (Cretaceous): ammonites, Texas, Mexico (41).
varves: Puente formation (Miocene), California (71).
Venezuela: echinoid, Eocene (21); mollusks, Cretaceous (72).
Ventricolaria (pelecypod): new name (33).
vertebrates: new Blancan locality (54).
Virginia: conifer, Triassic (7).
Welleraspis (trilobite): Cambrian (67).
Whitfield collection: types, University of California (61).
Wind River formation (Eocene): mammals, Wyoming (34).
Wyoming: Amsden formation, fauna and age (11); brachiopods, Permian (62); *Campylocynodon personi* (carnivore), Oligocene (12); mammals, Eocene (49); mammals, Lance formation (Cretaceous) (93); mammals, Wind River formation (Eocene) (34); ostracodes, Jurassic (63).